




Specialist Vascular Surgeons

**Supravalvular aortic arch stenosis:
A report of progressive disease and endovascular hybrid repair in a child.**

© 2025 Specialist Vascular Surgeons. All rights reserved.

This presentation is for informational purposes only and may not be copied, distributed, or used without written permission from the author.



Supravalvular aortic arch stenosis: A report of progressive disease and endovascular hybrid repair in a child.

Annals of Vascular Surgery – Brief Reports
and Innovations

Dr PP Mistry, et al.



Introduction

- SVAS – supraaortic stenosis
 - A disease of the arterial wall caused by reduced elastin (ELN)
- ELN arteriopathy
 - Usually results in thickening of the tunica intima and media, mostly affecting the ascending aorta.
 - Also reported in pulmonary and coronary arteries.
- SVAS may be non-syndromic or syndromic
 - Syndromic:
 - Complex genomic disorder that presents with neurobehavioural, craniofacial, cardiovascular and metabolic abnormalities.
 - Non-syndromic:
 - Caused by mutation of the ELN gene (auto-somal dominant inheritance)
 - ELN arteriopathy may result in discrete hourglass or diffuse long-segment stenosis of the supraaortic aorta.
 - Significant stenosis and high pressure gradients may result in hypertension, congestive cardiac failure, myocardial infarction and death.



Case Report

- 7 Year old male patient with complex, inherited, non-syndromic SVAS with numerous interventional procedures since birth.
 - Open surgical artery repair at 7 months for arch stenosis:
 - Bovine pericardial patch angioplasty of the ascending and descending aorta as well as the aortic arch via median sternotomy.
 - At age 7 he presented with refractory hypertension due to the SVAS with subsequent left ventricular hypertrophy and lower limb calf claudication.

Age-related anatomical abnormality, clinical presentation and intervention from birth to 7 years of age

Age	Anatomical abnormality	Clinical presentation	Surgical intervention
2-days	Patent foramen ovale (PPO) Peripheral pulmonary stenosis (PPS)	Ejection systolic murmur (ESM)	Nil
2-months	PPS	ESM	Nil
6-7-months	SVAS stenotic ascending aorta, arch, descending aorta, and PPS. Biventricular hypertrophy	Hypertension ESM	Pericardial patch / Bovine patch of ascending aorta / arch / proximal descending aorta
15-months	PPS	Hypertension	Balloon angioplasty PPS
17-months	Stenosis of LCCA and LSCA. Stenotic arch, proximal descending arch, LCCA, LSCA, PPS	Hypertension	Balloon angioplasty PPS Complication: pulmonary haemorrhage
27-months & 3-years	Angiogram: progressive narrowing of arch and proximal descending aorta	Progressive hypertension Anti-hypertensive escalated	Nil
7-years	Stenosis of the arch and proximal descending aorta Stenosis of the ostia of the LSCA and LCCA	Hypertension Left ventricular hypertrophy Lower limb claudication	Hybrid surgery

Pre-operative computed tomography (CT) angiogram

Findings:

- High-grade stenoses of the:
 - mid-arch
 - Proximal descending aorta
 - Left common carotid artery (LCCA)
 - Left subclavian artery (LSCA)
 - Pulmonary artery segmental branches
- The ascending aorta, brachiocephalic artery and right common carotid artery appeared free of stenoses.

Posterior oblique view of the aorta and great vessels showing high-grade stenosis of the proximal descending aortic (black arrow) and proximal LSCA high-grade stenosis (blue arrow)





Hybrid procedure – single stage

- Due to re-stenosis of the arch and ostial stenoses of the LCCA and LSCA, a hybrid procedure was done.
 - Extra-anatomical bypass:
 - Standard right to left CCA bypass and a RCCA to LSCA bypass using 6mm polytetrafluoroethylene (PTFE).
 - No carotid shunting used as the external carotid and vertebral arteries were adequate in maintaining cerebral perfusion.
 - Ligation of the stenosed proximal LCCA and LSCA.
 - Endovascular procedure:
 - Standard vascular access of the right femoral artery.
 - Angiography confirmed adequate cerebral perfusion.
 - Standard endovascular stent grafting from the brachiocephalic artery to the proximal descending aorta using an 11mmx100mm nitinol Gore Viabahn covered stent.
 - Residual stenosis of 50% (sustained pressure gradient of 40mmHg) following balloon dilatation.
 - Deployed a second 14x45mm stainless steel Z-Med stent graft.
 - The arch diameters were 10mm proximally and 4mm at the narrowest arch segment and 11mm at the mid-descending aorta.
 - The 2nd stent graft was placed as an inlay within the 1st stent graft.
 - A 10mm high pressure balloon was used to expand both stent grafts adequately.



Outcome:

Immediately post-operative

- No post-op complications.
- Improved calf claudication.
- His blood pressure appeared to improve to normal values for his age.
 - He was discharged with no anti-hypertensive medications.



Outcome: Follow-up

- 2 Months:
 - Normotensive → 108/47mmHg, normal range for height percentile = 113-116/72-77mmHg.
 - Anti-hypertensives introduced at age 6 were stopped after the procedure.
 - Claudication symptoms had resolved.
- 1 Year:
 - No clinical recurrence of hypertension or calf claudication
 - No further anti-hypertensives required.
 - Scripted Aspirin 75mg/daily.
 - No neurological symptoms reported.



CT angiogram follow-up


- 6 – Months:
 - Patent extra-anatomical bypasses and adequately expanded stent grafts.
- 1 Year:
 - High-grade stenosis at the ostium of the brachiocephalic artery.
 - Anastomotic stenoses of the previous bypass grafts.
 - The aortic stent grafts (SG-SG) remained relatively patent.

CT angiogram follow-up

Anterior posterior view showing the results of the hybrid procedure (black arrow) and progressive stenosis of the brachiocephalic ostium (yellow arrow).
AS = anastomotic stenosis.



Sagittal view after the hybrid procedure revealing patent stent grafts in the aortic arch.






Discussion

- Highlights the clinical consequences of ELN arteriopathy affecting the supraaortic aorta, the branches of the arch and pulmonary arterial branches
- Progressive arteriopathy:
 - Patch aortoplasty at 7 months (open surgery).
 - Restenosis of the arch at 7 years (hybrid surgery).
 - Multiple new ostial and anastomotic stenoses at 8 years (1 year follow up CT).



Discussion

- Open surgery is regarded as the gold standard for supraaortic coarctation, but reports indicate high rates of re-stenosis and re-operation.
 - Imamura et al (2010) Surgery for SVAS
 - 49 patients (age range 24 days to 24 years)
 - 25 years follow up
 - 86% underwent patch repair
 - 33% of re-operation due to stenosis.




Discussion

- **Shahanavaz et al (2020)** Balloon expandable stents on residual stenosis and hypertension
 - 57 patients (age range 4days – 42 years)
 - 7 Centres
 - 37 patients followed up
 - 8 patients required re-intervention.
- **Brown et al (2002)** Standard patch aortoplasty
 - 101 patients
 - 38 year follow up
 - 14% required re-operation at least once
- **Deo et al (2012)** Single-patch repair
 - 53 year follow up
 - 32 re-intervention procedures in 16 patients
 - Risk factors: young age at initial surgery and aortic stenosis.




Discussion

- Currently no comparative studies between endovascular intervention and open surgery in the treatment of non-syndromic SVAS patients.
- This case highlights the progressive stenotic complications associated with ELN arteriopathy.



Discussion

- Endovascular intervention may be better suited in high risk patients that have had multiple open surgical procedures.
- Adult endovascular interventionalists need to be aware of ELN arteriopathy and its progressive nature.
- Endovascular strategies may need to be modified for younger patients.
- In this case the stent-in-stent procedure allow for subsequent balloon angioplasty should it be needed.
- Proper endovascular planning and anticipation of progressive multi-vessel stenosis is important.



Conclusion

- Endovascular intervention using stent grafts may be considered an alternative to open surgery in aortic arch re-stenosis due to ELN arteriopathy.
- An extra-anatomical bypass may be necessary to facilitate stent graft placement.
- Although this procedure was feasible in this 7 year old and appeared to alleviate the symptoms, lifelong monitoring and possible subsequent intervention is mandatory.